

Observing solar-like oscillations

Timothy R. Bedding* and Hans Kjeldsen†

**School of Physics, University of Sydney, 2006, Australia*

†Danish AsteroSeismology Centre (DASC), University of Aarhus, Denmark

Abstract. The past few years have seen great progress in observing oscillations in solar-type stars, lying on or just above the main sequence. We review the most recent results, most of which were obtained using high-precision velocity measurements. We also briefly discuss observations of more evolved stars, namely G, K and M giants and supergiants.

Keywords: stellar oscillations

PACS: 97.10.Sj Pulsations, oscillations, and stellar seismology

MAIN-SEQUENCE AND SUBGIANT STARS

The past few years have seen great progress in observing oscillations in solar-type stars, lying on or just above the main sequence. Most of the recent results have come from high-precision Doppler measurements using spectrographs such as CORALIE, HARPS, UCLES and UVES. There are ambitious plans to build SONG (Stellar Oscillations Network Group), which will be a global network of small telescopes equipped with high-resolution spectrographs and dedicated to asteroseismology and planet searches (Grundahl et al., 2006).

Meanwhile, photometry from space is also making great progress. The WIRE and MOST missions have reported oscillations in several stars, although not without controversy, as discussed below. The recent launch and successful commissioning of COROT promises exciting new results (E. Michel et al., these Proceedings), as does the Kepler mission due for launch in early 2009.

Observations of solar-like oscillations are accumulating rapidly, and measurement have now been reported for several main-sequence and subgiant stars. The following list includes the most recent observations and is ordered according to decreasing stellar density (i.e., decreasing large frequency separation):

τ Cet (G8 V): this star was observed with HARPS by T. C. Teixeira et al. (in prep.)

70 Oph A (K0 V): this is the main component of a spectroscopic visual binary (the other component is K5 V). It was observed over 6 nights with HARPS by Carrier & Eggenberger (2006), who found $\Delta\nu = 162 \mu\text{Hz}$ but were not able to give unambiguous mode identifications from these single-site data.

α Cen B (K1 V): Oscillations in this star were detected from single-site observations with CORALIE by Carrier & Bourban (2003). Dual-site observations with UVES and UCLES (see Fig. 1) allowed measurement of nearly 40 modes and of the mode lifetime (Kjeldsen et al., 2005).




FIGURE 1. Power spectrum of α Cen B from velocity observations. Figure from Kjeldsen et al. (2005).




FIGURE 2. Four oscillation modes in α Cen A from the WIRE power spectrum, with fits that indicate the linewidth and rotational splitting. Figure from Fletcher et al. (2006).

α Cen A (G2 V): The clear detection of p-mode oscillations in α Cen A by Bouchy & Carrier (2002) using the CORALIE spectrograph represented a key moment in asteroseismology. This was followed by a dual-site campaign on this star with UVES and UCLES (Butler et al., 2004) that yielded more than 40 modes, with angular degrees of $l = 0$ to 3 (Bedding et al., 2004). The mode lifetime is about 2–4 days and there is now evidence of rotational splitting from photometry with the WIRE satellite analysed by Fletcher et al. (2006) (see Fig. 2) and also from ground-based spectroscopy with HARPS (Bazot et al., 2007).

μ Ara (G3 V): this star has multiple planets. Oscillations were measured over 8 nights using HARPS by Bouchy et al. (2005) (see Fig. 3) and the results were modelled by Bazot et al. (2005). They found $\Delta\nu = 90 \mu\text{Hz}$ and identified over 40 frequencies, with possible evidence for rotational splitting.

HD 49933 (F5 V): this is a primary target for the COROT space mission and was observed over 10 nights with HARPS by Mosser et al. (2005). They reported a surprisingly high level of velocity variability on timescales of a few days. This was also present as line-profile variations and is therefore presumably due to stellar activity. The observations showed excess power from p-mode oscillations and the authors determined the large separation ($\Delta\nu = 89 \mu\text{Hz}$) but were not able to extract




FIGURE 3. Radial velocity time series of the star μ Ara made over 8 nights with the HARPS spectrograph. Figure from Bouchy et al. (2005).

individual frequencies.

β Vir (F9 V): oscillations in this star were detected in a weather-affected two-site campaign with ELODIE and FEROS by Martić et al. (2004). Subsequently, Carrier et al. (2005) used CORALIE with good weather but a single site, and reported 31 individual frequencies. Those results were modelled by Eggenberger & Carrier (2006), who also reported tentative evidence for rotational splittings. The large separation is $72 \mu\text{Hz}$.

Procyon A (F5 IV): At least eight separate velocity studies have reported an excess in the power spectrum, beginning with that by Brown et al. (1991). The most recent examples were reported by Martić et al. (2004), Eggenberger et al. (2004), Bouchy et al. (2004) and Leccia et al. (2007). These studies agreed on the location of the excess power (around $0.5\text{--}1.5 \text{ mHz}$) but they disagreed on the individual oscillation frequencies. However, a consensus has emerged that the large separation is about $55 \mu\text{Hz}$.

Controversy was generated when photometric observations obtained with the MOST satellite by Matthews et al. (2004) failed to reveal evidence for oscillations. However, Bedding et al. (2005) argued that the MOST non-detection was consistent with the ground-based data and Régulo & Roca Cortés (2005) suggested that the signature of oscillations is indeed present in the MOST data at a low level. Using space-based photometry with the WIRE satellite, Bruntt et al. (2005)




FIGURE 4. Échelle diagram of oscillation frequencies in β Hyi. Figure from Bedding et al. (2007).

extracted parameters for the stellar granulation and found evidence for an excess due to p-mode oscillations.

A multi-site spectroscopic campaign on Procyon was carried out in January 2007, which has confirmed that this star does indeed show solar-like oscillations (Arentoft et al. in prep.; Bedding et al., in prep.).

β Hyi (G2 IV): In 2005 this star was the target for a two-site campaign using HARPS and UCLES Bedding et al. (2007), which resulted in the clear detection of mixed modes (see Fig. 4). The large separation is $57.5 \mu\text{Hz}$. Combining this value with the angular diameter measured using interferometry gave the stellar mass to an accuracy of 2.7% (North et al., 2007). This is probably the most precise mass determination of a solar-type star that is not in a binary system, illustrating the power of combining asteroseismology and interferometry (see also Creevey et al. 2007 and references therein).

δ Eri (K0 IV): Carrier et al. (2003) observed this star over 12 nights in 2001 with CORALIE and found a large separation of $44 \mu\text{Hz}$.

η Boo (G0 IV): The claimed detection of oscillations by Kjeldsen et al. (1995), based on fluctuations in Balmer-line equivalent-widths, has now been confirmed by further equivalent-width and velocity measurements by the same group (Kjeldsen et al., 2003) and also by independent velocity measurements with the CORALIE spectrograph (Carrier et al., 2005). With the benefit of hindsight, we can now say that η Boo was the first star for which the large separation and individual frequencies were measured. However, there is still disagreement on some of the individual frequencies, which reflects the subjective way in which genuine oscillation modes must be chosen from noise peaks and corrected for daily aliases. Fortunately, the large separation is $\Delta\nu=40 \mu\text{Hz}$, which is half way between integral multiples of the $11.57\text{-}\mu\text{Hz}$ daily splitting ($40/11.57 = 3.5$). Even so, daily aliases are problematic, especially because some of the modes in η Boo appear to

be shifted by avoided crossings.

Spaced-based observations of η Boo, made with the MOST satellite, have generated considerable controversy. Guenther et al. (2005) showed an amplitude spectrum (their Fig. 1) that rises towards low frequencies in a fashion that is typical of noise from instrumental and stellar sources. However, they assessed the significance of individual peaks by their strength relative to a fixed horizontal threshold, which naturally led them to assign high significance to peaks at low frequency. They did find a few peaks around $600 \mu\text{Hz}$ that agreed with the ground-based data, but they also identified eight of the many peaks at much lower frequency ($130\text{--}500 \mu\text{Hz}$), in the region of rising power, as being due to low-overtone p-modes. Those peaks do line up quite well with the regular $40 \mu\text{Hz}$ spacing, but extreme caution is needed before these peaks are accepted as genuine. This is especially true given that the orbital frequency of the spacecraft ($164.3 \mu\text{Hz}$) is, by bad luck, close to four times the large separation of η Boo ($164.3/40 = 4.1$). Models of η Boo based on the combination of MOST and ground-based frequencies have been made by Straka et al. (2006).

ν Ind (G0 IV): this a metal-poor subgiant ($[\text{Fe}/\text{H}] = -1.4$) which was observed from two sites using UCLES and CORALIE (Bedding et al., 2006; Carrier et al., 2007). The large separation of $24 \mu\text{Hz}$, combined with the position of the star in the H-R diagram, indicated that the star has a low mass ($0.85 \pm 0.04 M_{\odot}$) and is at least 9 Gyr old (see Fig. 5).

G AND K GIANTS

Oscillations in a few G and K giant stars have been detected using ground-based velocity observations, indicating periods of a few to several hours. Oscillations in ξ Hya (G7 III) (Frandsen et al., 2002; Stello et al., 2004) measured using CORALIE appear to have mode lifetimes of only about 2 days. The CORALIE and ELODIE spectrographs have been used to find excess power and a possible large separation for both ε Oph (G9 III) (De Ridder et al., 2006) and η Ser (K0 III) (Barban et al., 2004). Hekker et al. (2006) have analysed the line-profile variations and found evidence for non-radial oscillations in some of these stars. Most recently, Hatzes & Zechmeister (2007) reported evidence for oscillations in the K giant star β Gem, although this was based on a rather short time series (20 hours).

Spaced-based photometry of α UMa (K0 III) with WIRE produced evidence for multi-mode oscillations (Buzasi et al., 2000). More recently, photometry of ε Oph with MOST confirmed the ground-based detection of oscillations, establishing the large separation to be $5.1 \mu\text{Hz}$ and indicating a mode lifetime of 2–3 d (Barban et al., 2007). Mode lifetimes of a few days are the same as those in the Sun but these giant stars oscillate about 50 times more slowly. Such short lifetimes, if confirmed as typical, would significantly limit the prospects for asteroseismology on red giants.

Photometry of clusters is another way to search for oscillations in K giants. Edmonds & Gilliland (1996) observed K giants in the globular cluster 47 Tuc using the *Hubble Space Telescope* over 38.5 hr and found variables with periods of 2–4 days and semi-amplitudes of 5–15 mmag. More recently, multi-site ground-based




FIGURE 5. Parameters of ν Ind for three different choices of α , the mixing-length parameter. The thin error bars show the range of each parameter based on classical measurements alone (luminosity and temperature), while the thick bars include the constraint provided by the large frequency separation. The dashed line at 13.7 Gyr indicates the upper limit set by age of the universe from cosmology. Figure from Bedding et al. (2006).

observations have given tentative evidence for oscillations in two open clusters, namely M67 (Stello et al., 2007) and M4 (Frandsen et al., 2007).

Finally, we note that photometry of Arcturus (K1.5 III) has been reported by Tarrant et al. (2007) from 2.5 yr of observations made by the Solar Mass Ejection Imager (SMEI) on board the Coriolis satellite. The observations indicate a strong mode at $3.5\mu\text{Hz}$ with a damping time of 24 d. This mode lifetime is also short when we take into account the period of the oscillations (3.3 d).

M GIANTS AND SUPERGIANTS

If we define solar-like oscillations to be those excited and damped by convection then we might expect to see such oscillations in all stars on the cool side of the instability strip. Evidence for solar-like oscillations in semiregular variables, based on visual ob-




FIGURE 6. Power spectra of red supergiants from visual observations (thin lines) with Lorentzian fits (thick lines). Figure from Kiss et al. (2006).

servations by groups such as the AAVSO, has already been reported. This was based on the amplitude variability of these stars (Christensen-Dalsgaard et al., 2001) and on the Lorentzian profiles of the power spectra (Bedding et al., 2005).

Recently, Kiss et al. (2006) used visual observations from the AAVSO to show that red supergiants such as Betelgeuse (α Ori), which have masses of $10\text{--}30M_{\odot}$, also have Lorentzian profiles in their power spectra (see Fig. 6).

ACKNOWLEDGMENTS

This work was supported financially by the Australian Research Council, the Science Foundation for Physics at the University of Sydney, and the Danish Natural Science Research Council.

REFERENCES

- Barban C., de Ridder J., Mazumdar A., et al., 2004, in Danesy D., ed., SOHO 14/GONG 2004 Workshop, Helio- and Asteroseismology: Towards a Golden Future ESA SP-559. p. 113
- Barban C., Matthews J. M., de Ridder J., et al., 2007, A&A, 468, 1033
- Bazot M., Bouchy F., Kjeldsen H., Charpinet S., Laymand M., Vauclair S., 2007, A&A, 470, 295
- Bazot M., Vauclair S., Bouchy F., Santos N. C., 2005, A&A, 440, 615
- Bedding T. R., Butler R. P., Carrier F., et al., 2006, ApJ, 647, 558
- Bedding T. R., Kiss L. L., Kjeldsen H., et al., 2005, MNRAS, 361, 1375
- Bedding T. R., Kjeldsen H., Arentoft T., et al., 2007, ApJ, 663, 1315
- Bedding T. R., Kjeldsen H., Bouchy F., et al., 2005, A&A, 432, L43
- Bedding T. R., Kjeldsen H., Butler R. P., et al., 2004, ApJ, 614, 380
- Bouchy F., Bazot M., Santos N. C., Vauclair S., Sosnowska D., 2005, A&A, 440, 609
- Bouchy F., Carrier F., 2002, A&A, 390, 205
- Bouchy F., Maeder A., Mayor M., Mégevand D., Pepe F., Sosnowska D., 2004, Nat, 432, 7015
- Brown T. M., Gilliland R. L., Noyes R. W., Ramsey L. W., 1991, ApJ, 368, 599
- Bruntt H., Kjeldsen H., Buzasi D. L., Bedding T. R., 2005, ApJ, 633, 440
- Butler R. P., Bedding T. R., Kjeldsen H., et al., 2004, ApJ, 600, L75
- Buzasi D. L., Catanzarite J., Laher R., et al., 2000, ApJ, 532, L133
- Carrier F., Bouchy F., Eggenberger P., 2003, in Thompson M. J., Cunha M. S., Monteiro M. J. P. F. G., eds, Asteroseismology Across the HR Diagram Kluwer, p. P311
- Carrier F., Bourban G., 2003, A&A, 406, L23
- Carrier F., Eggenberger P., 2006, A&A, 450, 695
- Carrier F., Eggenberger P., Bouchy F., 2005, A&A, 434, 1085
- Carrier F., Eggenberger P., D'Alessandro A., Weber L., 2005, NewA, 10, 315
- Carrier F., Kjeldsen H., Bedding T. R., et al., 2007, A&A, 470, 1059
- Christensen-Dalsgaard J., Kjeldsen H., Mattei J. A., 2001, ApJ, 562, L141
- Creevey O. L., Monteiro M. J. P. F. G., Metcalfe T. S., et al., 2007, ApJ, 659, 616
- De Ridder J., Barban C., Carrier F., et al., 2006, A&A, 448, 689
- Edmonds P. D., Gilliland R. L., 1996, ApJ, 464, L157
- Eggenberger P., Carrier F., 2006, A&A, 449, 293
- Eggenberger P., Carrier F., Bouchy F., Blecha A., 2004, A&A, 422, 247
- Fletcher S. T., Chaplin W. J., Elsworth Y., Schou J., Buzasi D., 2006, MNRAS, 371, 935
- Frandsen S., Bruntt H., Grundahl F., et al., 2007, A&A, in press
- Frandsen S., Carrier F., Aerts C., et al., 2002, A&A, 394, L5
- Grundahl F., Kjeldsen H., Frandsen S., et al., 2006, Mem. Soc. Astron. Ital., 77, 458
- Guenther D. B., Kallinger T., Reegen P., et al., 2005, ApJ, 635, 547
- Hatzes A. P., Zechmeister M., 2007, Astrophys. J., Lett., in press, [arXiv:astro-ph/0709.1406](https://arxiv.org/abs/0709.1406)
- Hekker S., Aerts C., De Ridder J., Carrier F., 2006, A&A, 458, 931
- Kiss L. L., Szabo G. M., Bedding T. R., 2006, MNRAS, 372, 1721
- Kjeldsen H., Bedding T. R., Baldry I. K., et al., 2003, AJ, 126, 1483
- Kjeldsen H., Bedding T. R., Butler R. P., et al., 2005, ApJ, 635, 1281
- Kjeldsen H., Bedding T. R., Viskum M., Frandsen S., 1995, AJ, 109, 1313
- Leccia S., Kjeldsen H., Bonanno A., Claudi R. U., Ventura R., Paternò L., 2007, A&A, 464, 1059
- Martić M., Lebrun J.-C., Appourchaux T., Korzennik S. G., 2004, A&A, 418, 295
- Martić M., Lebrun J. C., Appourchaux T., Schmitt J., 2004, in Danesy D., ed., SOHO 14/GONG 2004 Workshop, Helio- and Asteroseismology: Towards a Golden Future ESA SP-559. p. 563
- Matthews J. M., Kuschnig R., Guenther D. B., et al., 2004, Nat, 430, 51, Erratum: 430, 921
- Mosser B., Bouchy F., Catala C., et al., 2005, A&A, 431, L13
- North J. R., Davis J., Bedding T. R., et al., 2007, MNRAS, 380, L83
- Régulo C., Roca Cortés T., 2005, A&A, 444, L5
- Stello D., Bruntt H., Kjeldsen H., Bedding T. R., Arentoft T., et al., 2007, MNRAS, 377, 584
- Stello D., Kjeldsen H., Bedding T. R., et al., 2004, Sol. Phys., 220, 207
- Straka C. W., Demarque P., Guenther D. B., Li L., Robinson F. J., 2006, ApJ, 636, 1078
- Tarrant N. J., Chaplin W. J., Elsworth Y., et al., 2007, MNRAS, in press, [arXiv:astro-ph/0706.3346](https://arxiv.org/abs/0706.3346)